

Extract of metabarcoding.org

<http://metabarcoding.org/spip.php?article75>

# **ECOGEN - Ecosystem change and species persistence over time: a genome-based approach**

- Position opportunities -

Publication date: Tuesday 9 February 2016

---

**Copyright © metabarcoding.org - All rights reserved**

---

## Main goal

---

**Develop high taxonomic resolution ancient environmental DNA methods in order to evaluate how drivers of change (human, climate, biota) affect species persistence and ecosystem tipping points in arctic-alpine biomes**

## Subgoals

---

- 1 - Improve methods for full genome analyses of environmental DNA**
- 2 - Compile palaeo data to plan a balanced design of climatic and human impacts**
- 3 - Do full genome analyses of lake cores to obtain information on past presence and abundance of vascular plant species and key herbivores**
- 4 - Identify biotic drivers and disentangle their effects from human land use and climate change on ecosystem resilience and ecosystem services**
- 5 - Estimate species persistence across periods of changes and identify factors causing extinction**
- 6 - Provide methods and knowledge to inform species conservation and ecosystem management**

## Project tasks

---

We are seeking a research fellow that will join a cross-disciplinary team (ecology, palaeoecology, genetics, archaeology, geology, niche modelling) working in two geographical regions (Norway and the Alps). We will expand our knowledge on past vascular plant and animal diversity and abundance at a taxonomic depth that has not been possible until now due to methodological limitations. We expect to disentangle the effects of past human land-use (hunting, husbandry, burning, agriculture), climate change, and biota on species and ecosystem changes and thereby be able to answer questions central to our understanding of our biological resources, such as the level of persistence of species and resilience of ecosystems to environmental drivers, the extinction risk of species, and the capacity of mountain landscape to buffer against these changes. By identifying drivers of shifts in ecosystem services through time, we may inform future management.

The successful candidate will work on development of improved technology. The most recent applications of ancient plant DNA analyses are largely developed by our team (Taberlet et al. 2007, Sønstedt et al. 2010, Yoccoz et al. 2012, Willerslev et al. 2014). For vascular plants, the 50-100 base pair long P6 loop region of the chloroplast trnL (UAA) intron is used in a PCR based method (Taberlet et al. 2007), which allows identification of all plant families, most genera (>75%), and one third of the species (Sønstedt et al. 2010). When applied to modern lake sediments, half of the species present within 2 m of lakes were detected (Alsos et al. In prep.). Our recent study of an 8500 year old core from Svalbard show that the method detect all except two genera identified in a macrofossil study from the same lake (Alsos et al. 2015). Overall, 1.2 times more taxa of vascular plants were identified with ancient DNA than macrofossils, and the number of taxa identified per sample was 2.7 times higher for the former. Thus, DNA analyses of Holocene lake sediment can reveal the presence of rare taxa and thereby allow for a better estimation of species persistence. Further, the method is now resource-efficient and repeatable, and it can be extended to any group of organisms, given that a DNA reference library and adequate primers are developed. It also allows for semi-quantitative interpretations based on the number of PCR repeats where taxa are identified (Pansu et al. 2015). We will do minor optimization of these applications and analyse all samples for vascular plants and key herbivores.

Contact : Prof. Inger Greve Alsos  
Tromsø University Museum  
NO-9037 Tromsø  
Norway  
Email: [inger.g.alsos@uit.no](mailto:inger.g.alsos@uit.no)